



Review

Neuromuscular Electrical Stimulation for Dysphonia and Dysarthria: A Systematic Review

Franziska Schreiber^{1,2,*#}, Nayara de Lima Froio^{2,3,#}, Roser Coll-Fernández^{2,4,#}, Fernanda Queiroz Petrus^{2,5}, Bethânia Soares dos Santos^{2,6,7}, Angélica Flores Flores^{2,8}, Luis A. Morales Ojeda^{2,9}, Laura de Leon Benedetti^{2,10}, Vinicius Trindade^{2,11,12}, Mustafa Reha Dodurgali², Claudia Renata dos Santos Barros^{2,13}, Bianca Barros Parron Fernandes^{2,4}, Christiany M. Tapia Argudo^{2,14}, Cristina Stephan^{2,15} and Francesca Buffone^{2,16, 17}

¹ Institute of Pathology, Faculty of Medicine, Ludwig-Maximilians University Munich, Thalkirchner Straße 36, 80337 Munich, Germany;

² PPCR, Harvard T. H. Chan School of Public Health – ECPE, Boston, MA, USA

Rest of author's affiliation at the end of the manuscript.

* Correspondence: Franziska Schreiber, franziska.schreiber13@gmail.com

These authors have contributed equally to this work

Received: 11/09/2022; 12/23/2022; published: 04/04/23.

ABSTRACT:

Introduction: People with speech disorders such as dysarthria and dysphonia have difficulty producing speech, which is often related to speech muscle dysfunction. Neuromuscular electrical stimulation (NMES) can activate motor units and improve their tropism and functionality. Therefore, this systematic review aimed to evaluate the effects of NMES on the phonation, voice quality, intelligibility, speech muscle strength, and quality of life of patients with dysarthria and dysphonia.

Methods: This systematic review was performed according to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses guidelines. In September 2022, we searched the EMBASE, PubMed, and Cochrane databases for studies that evaluated the effect of NMES on dysarthria and dysphonia. Randomized controlled trials, observational studies, case series, and case reports were included in the search strategy. The risk of bias in each study was assessed using the ROBINS-I tool.

Results: This review included four studies designed as case studies or non-randomized trials with a low risk of bias. NMES showed a positive effect on voice quality and phonation in patients with dysarthria and/or dysphonia. Reported improvements included restored function of the muscles responsible for laryngeal elevation, a reduction in voice breathiness and strain, and enhanced articulation.

Discussion: NMES seems to be a promising tool for the rehabilitation of patients with speech disorders. Randomized clinical trials should be conducted to confirm these benefits.

Keywords: Neuromuscular stimulation; peripheral electrical stimulation; NMES; electrical stimulation; dysarthria; voice disorders; articulation disorders; dysphonia; phonation disorder; voice disturbance

Academic Editor: Felipe Fregni

Peer-reviewers: Augusto J Mendes; Salvatore Minnella; Nadine Aranis;

Copyright: © 2022 by the authors. Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).



DOI: <http://dx.doi.org/10.21801/ppcrj.2022.84.10>

Introduction

People with speech disorders have difficulty producing speech, which is often related to speech muscle dysfunction. Dysarthria and dysphonia, among the most prevalent speech disorders, result from various pathologies, such as vocal fold paralysis, spasmodic dysphonia, traumatic brain injury, post-surgical meningioma, meningitis, stroke, and other neurologic diseases (Sellars et al., 2005; Almeida et al., 2022). Patients with these conditions face considerable barriers in their activities and participation in social and civil life (Wray et al., 2019; Brady et al., 2011). Therefore, improving dysarthria and dysphonia is important for rehabilitating affected patients.

Neuromuscular electrical stimulation (NMES) involves intermittent peripheral nerve stimulation that activates the motor units and improves their trophism and functionality. Furthermore, it promotes modulation of the sensorimotor cortex and spinal motor neurons (Maffiuletti, 2010). Since the early 1960s, NMES has been used to aid post-stroke recovery (Ijzerman et al., 2009), and several studies have proven its benefits in patients with post-stroke dysphagia (Oh et al., 2017; Oh et al., 2020; Alamer et al., 2020; Park et al., 2016). Since the muscles involved in dysphagia and those involved in dysarthria/dysphonia show an overlap, it seems reasonable to use NMES to improve speech disorders. However, to date, there are no evidence-based recommendations for NMES implementation in dysarthria/dysphonia rehabilitation.

This systematic review makes a novel contribution to the literature by evaluating the effects of NMES on the phonation, voice quality, intelligibility, speech muscle strength, and quality of life of patients with dysarthria and dysphonia and comparing these effects with those of other speech therapies, in the short and long term.

Materials and Methods

Study design

The Preferred Reporting Items for Systematic Reviews and Meta-Analyses guidelines were followed in this study (Page et al., 2021).

Search strategy

This is a systematic review of NMES for treating patients with dysarthria and/or dysphonia. In September 2022, we systematically searched the electronic databases EMBASE, PubMed, and Cochrane for articles published in English, Spanish, Portuguese, and German using the search terms ("dysphonia'[MeSH

Terms]'" OR "Dysphonia OR 'Hyperkinetic Dysphonia' OR 'Neurologic Adductor Spastic Dysphonia' OR 'Organic Tremor Dysphonia' OR 'Phonation Disorder*' OR 'Spastic Dysphonia'") OR ("dysarthria[MeSH Terms]" OR "(Dysarthria OR 'Dysarthros*' OR 'Dysarthrias' OR 'Flaccid Dysarthria*' OR 'Guttural Dysarthria*' OR 'Mixed Dysarthria*' OR 'Scanning Dysarthria*' OR 'Spastic Dysarthria*') AND ("electric stimulation therapy[MeSH Terms]" OR "'Electric Stimulation Therapy' OR 'Electric* Stimulation Therapy' OR 'Electrotherapy' OR 'Interferential Current Electrotherapy' OR 'Stimulation Therapy, Electrical' OR 'Therapeutic Electric* Stimulation' OR 'NMES' OR 'Electrical stimulation*' OR 'Neuromuscular stimulation'").

Eligibility criteria

We included studies that evaluated the effects of NMES on dysarthria and dysphonia. The outcomes were changes in quantitative or qualitative measurements of phonation, voice quality, intelligibility, speech muscle strength, and quality of life (for instance: assessment of articulation, vowel distortion, hypernasality, loudness and pitch variation). We included randomized controlled trials (RCTs), observational studies, case series, and case reports, published in the previous 10 years. We excluded studies on patients with speech or language disorders other than dysarthria and dysphonia, such as aphasia and speech apraxia. Those involving transcutaneous electrical nerve stimulation or deep brain stimulation were also excluded.

Study selection

The electronic database search was conducted by one reviewer. After duplicates were removed, titles and abstracts were independently evaluated by three reviewers based on the eligibility criteria. Thereafter, eight reviewers screened the full text of each potential article. Disagreements were settled through group discussion among the eight reviewers to reach a consensus about each article.

Assessment of risk of bias in individual studies

Two reviewers independently assessed the risk of bias for each of the four papers that were finally included using the ROBINS-I tool for non-randomized trials (Sterne et al., 2016).


Data extraction and analysis

Articles found in the databases were exported to the Rayyan AI platform. We followed the PICOS criteria for data extraction as follows: (P) Population, adult patients with dysphonia and/or dysarthria; (I) Intervention, NMES; (C) Comparison, any other treatment, including placebo; (O) Outcome, changes in quantitative or qualitative measurements of phonation, voice quality, intelligibility, speech muscle strength, and quality of life; (S) Studies, RCTs, observational studies, case series and case reports.

Results

The systematic search yielded 542 articles, with 81 duplicated studies. After screening the titles and abstracts, 15 papers remained. After reading the entire papers, we excluded 11 articles for the following reasons: patients with diagnoses other than dysarthria/dysphonia (n=5), interventions other than NMES (n= 1), publication date outside the stipulated time frame (n= 2), language other than those stipulated (n=1), and conference abstracts (n=2). Finally, four articles were included in this review (Figure 1). The details of the included studies are shown in Table 1. All the included studies were classified as having a low risk of bias according to the ROBINS-I guidelines (Sterne et al., 2016).

Figure 1. Preferred Reporting Items for Systematic Reviews and Meta-Analyses flow diagram



Effect of NMES on voice quality

Ko et al. (2016) examined the impact of laryngopharyngeal NMES therapy in patients with stroke or traumatic brain injury who had speech disorder and dysphagia. Eighteen patients were evaluated, of whom only six received conventional swallowing training and twelve received this training plus NMES. The authors reported an improvement in vocal fold vibration/tension and restoration in the function of the muscles responsible for laryngeal elevation.

In a retrospective case study, Guzman et al. (2014) reported positive effects of NMES therapy in addition to traditional vocal exercises on the voice quality of two female patients with superior laryngeal nerve weakness. With the addition of NMES, a significant improvement in voice breaking and vocal range, a decrease in voice breathiness, and an increase in voice resonance could be attained.

Safi et al. (2018) included two patients with mildly to moderately dysarthric speech after a cerebrovascular accident in a study with a single-subject design that involved a combination of oral motor exercise with NMES. Submental NMES showed no effect on lingual strength, but facial NMES led to an improvement of labial strength in both patients; this can ultimately contribute to an improvement in voice quality, considering the complexity and involvement of different muscles for speech production.

Furthermore, NMES showed promising results in one patient with severe dysarthria. Berenati et al. (2021) describe the strengthening of muscle contractions and improvement in vocal strain in an 18-year-old with severe speech impairment after post-anoxic brain injury.

Effects of NMES on phonation

Ko et al. (2016) report positive effects on phonation after 2 and 4 weeks of NMES therapy in patients with stroke. Additionally, Guzman et al. (2014) describe an improvement in soft phonation index values after 17 and 8 sessions with NMES, respectively.

Furthermore, as reported by Berenati et al. (2021), NMES can improve articulation and hypernasality, contributing to improved phonation. The patient presented in the case study by Berenati et al. (2021) was unable to perform conventional speech therapy due to his impaired cognitive abilities. Consequently, his dysarthria severity was reduced solely through NMES therapy. These findings are especially remarkable in light of the reported increased amplitude

of the patient's trigeminal-facial reflex responses, which can be attributed to the induction of neuroplastic changes in the sensorimotor areas responsible for articulatory functions.

Effect of NMES on quality of life

Dysarthria and dysphonia impact patients' social life and well-being, and affect patients face restrictions in daily activities that require the use of their voice. In addition to the impact on patients' communicative quality of life, other aspects, such as psychological or social domains, may be impacted (Ma & Yiu, 2001). Despite this, Guzman et al. (2014), Ko et al. (2016), and Safi et al. (2018) acknowledge that this patient group can have a significantly decreased quality of life; however, none of the studies included in this review investigated this knowledge gap further.

Discussion

Laryngeal muscle weakness can result in vocal fold instability. Acoustic evaluations reveal that patients with stroke-related dysarthria have significant differences in frequency, jitter, shimmer, and noise-to-harmonic ratio compared with those of neurologically healthy controls (Wang et al., 2010). As a result, treatments that strengthen these muscles seem to be promising for improving speech disorders.

Laryngopharyngeal NMES can restore the function of muscles controlling vocal fold vibration, which can strengthen the muscles attached to the hyoid bone (namely, the mylohyoid and thyrohyoid muscles) and increase laryngeal elevation. Consequently, excessive quavering during speech can be reduced. Furthermore, NMES can improve the sound pressure level by strengthening vocal ligament and thyroarytenoid muscle tension, thereby increasing glottal resistance (McHenry & Reich, 1985).

Despite the possible benefits, we would like to highlight some limitations of the present studies. None of them was a RCT, thus the patients were not blinded, and the sample size was small. The maximum duration of NMES treatment was 4 weeks and maybe this might be too short to generate a significant and lasting effect (especially if we consider that an improvement in muscle function or neuromodulation is proportional to the frequency of the stimuli). In addition, 3 of the 4 studies did not include a comparison group.

Conclusions

In conclusion, NMES may be an effective and promising tool for the rehabilitation of patients with

speech disorders. The limited number of studies regarding its use for treating dysphonia and dysarthria in adults did not allow us to perform a meta-analysis and obtain statistically significant results supporting the benefit of the use of electrical stimulation in these patients.

To the best of our knowledge, no RCTs investigated this topic up to date. Nevertheless, to confirm the benefits of NMES in the treatment of dysphonia and dysarthria and to further explore the findings presented in this systematic review, conducting an RCT would be highly beneficial.

None of the present studies has evaluated the patients in the medium to long term, and thus the duration of the effects of NMES is not yet clear. We suggest this assessment in future studies. Furthermore, we encourage the evaluation of quality of life to highlight the benefits of NMES therapy. This could broaden the understanding of the effects of NMES on dysphonia and dysarthria and its impact on the patient's social life and well-being.

Funding: This research received no external funding

Acknowledgments: We would like to express our gratitude to our teaching assistants Caio Cesar dos Santos Kasai and Mariana Gasparoto for guiding us throughout this work. We wish to thank the PPCR team (faculty members, coordinators, reviewers) including Professor Felipe Fregni and Alma Sanchez Jimenez for their invaluable support.

Conflicts of Interest: Cristina Stephan is a medical affairs consultant for different pharmaceutical companies. All other authors declare no conflict of interest.

Authors' affiliations:

- ¹ *Institute of Pathology, Faculty of Medicine, Ludwig-Maximilians University Munich, Thalkirchner Straße 36, 80337 Munich, Germany;*
- ² *PPCR, Harvard T.H. Chan School of Public Health - ECPE, Boston, MA 02115, USA;*
- ³ *Hospital Alemão Oswaldo Cruz, R. Treze de Maio, 1815 - Bela Vista, São Paulo - SP, 01323-020, Brazil;*
- ⁴ *Rehabilitation Department, Hospital Universitari Germans Trias i Pujol, Carretera del Canyet s/n, 08916 Badalona, Spain;*
- ⁵ *School of Medicine, Pontifical Catholic University of Paraná (PUCPR), Av. Jockey Club, 485, Hípica, Londrina, PR, Brazil;*
- ⁶ *D'Or Institute Research and Education (IDOR) - Rio de Janeiro, 22281-100, Brazil;*
- ⁷ *Department of Clinical Research, Brazilian National Cancer Institute (INCA) - Rio de Janeiro, 20560-121, Brazil;*
- ⁸ *Laboratory of Immunopharmacology, Instituto Nacional de Enfermedades Respiratorias "Ismael Cosío Villegas", Calz. de Tlalpan 4502, Belisario Domínguez Secc 16, Tlalpan, 14080 Ciudad de México, CDMX;*

- ⁹ University of Southern California (USC), Institute of Urology, Keck School of Medicine 1975 Zonal Ave. Los Angeles, CA, 90033;
- ¹⁰ Department of Radiology, Children's hospital of Philadelphia, Philadelphia, USA;
- ¹¹ University of São Paulo Medical School, Department of Neurology, Division of Neurosurgery- 255 Dr Eneas de Carvalho Aguiar st, Pinheiros, Sao Paulo, SP, Brazil;
- ¹² AC Camargo Cancer Center, Rua Tamandaré, 753 - Liberdade, Sao Paulo - SP, Brazil, ZIP Code 01525-001;
- ¹³ Butantan Institute, 210 Universidade avenue, Sao Paulo - Brazil - CEP 05508-040;
- ¹⁴ DOCS Urgent Care, Orange, Connecticut, 06477. USA;
- ¹⁵ Step Medical Affairs Consulting, Barata Ribeiro st, Bela Vista, São Paulo, Brazil, Zip Code: 01308-000;
- ¹⁶ Division of Paediatric, Manima Non-Profit Organization Social Assistance and Healthcare, 20125 Milan, Italy;
- ¹⁷ Department of Research, SOMA—Istituto Osteopatia Milano, 20126 Milan, Italy;

References

- Alamer, A., Melese, H., & Nigussie, F. (2020). Effectiveness of Neuromuscular Electrical Stimulation on Post-Stroke Dysphagia: A Systematic Review of Randomized Controlled Trials. *Clinical Interventions in Aging*, Volume 15, 1521–1531. <https://doi.org/10.2147/CIA.S262596>
- Almeida, A. N. S. de, Cunha, D. A. da, Ferreira, S. L. de S., Guimarães, B. T. de L., Balata, P. M. M., & Silva, H. J. da. (2022). Effect of Electrical Stimulation on the Treatment of Dysphonia: A Systematic Review. *Journal of Voice*, 36(5), 650–660. <https://doi.org/10.1016/j.jvoice.2020.08.002>
- Berenati, M., Naro, A., Calabrò, C., Torrisi, M., Cardali, S. M., & Calabrò, R. S. (2021). Is Neuromuscular Electrical Stimulation Effective in Treating Severe Dysarthria: Insights from a Case Study. *Innovations in Clinical Neuroscience*, 18(10–12), 23–25.
- Brady, M. C., Clark, A. M., Dickson, S., Paton, G., & Barbour, R. S. (2011). The impact of stroke-related dysarthria on social participation and implications for rehabilitation. *Disability and Rehabilitation*, 33(3), 178–186. <https://doi.org/10.3109/09638288.2010.517897>
- Guzman, M., Rubin, A., Cox, P., Landini, F., & Jackson-Menaldi, C. (2014). Neuromuscular Electrical Stimulation of the Cricothyroid Muscle in Patients With Suspected Superior Laryngeal Nerve Weakness. *Journal of Voice*, 28(2), 216–225. <https://doi.org/10.1016/j.jvoice.2013.09.003>
- Ijzerman, M. J., Renzenbrink, G. J., & Geurts, A. C. (2009). Neuromuscular stimulation after stroke: From technology to clinical deployment. *Expert Review of Neurotherapeutics*, 9(4), 541–552. <https://doi.org/10.1586/ern.09.6>
- Ko, K. R., Park, H. J., Hyun, J. K., Seo, I.-H., & Kim, T. U. (2016). Effect of Laryngopharyngeal Neuromuscular Electrical Stimulation on Dysphonia Accompanied by Dysphagia in Post-stroke and Traumatic Brain Injury Patients: A Pilot Study. *Annals of Rehabilitation Medicine*, 40(4), 600–610. <https://doi.org/10.5535/arm.2016.40.4.600>
- Ma, E. P., & Yiu, E. M. (2001). Voice activity and participation profile: assessing the impact of voice disorders on daily activities. *Journal of speech, language, and hearing research: JSLHR*, 44(3), 511–524. [https://doi.org/10.1044/1092-4388\(2001/040\)](https://doi.org/10.1044/1092-4388(2001/040))
- Maffiuletti, N. A. (2010). Physiological and methodological considerations for the use of neuromuscular electrical stimulation. *European Journal of Applied Physiology*, 110(2), 223–234. <https://doi.org/10.1007/s00421-010-1502-y>
- McHenry, M. A., & Reich, A. R. (1985). Effective Airway Resistance and Vocal Sound Pressure Level in Cheerleaders with a History of Dysphonic Episodes. *Folia Phoniatrica et Logopaedica*, 37(5–6), 223–231. <https://doi.org/10.1159/000265802>
- Oh, D.-H., Park, J.-S., Kim, H.-J., Chang, M.-Y., & Hwang, N.-K. (2020). The effect of neuromuscular electrical stimulation with different electrode positions on swallowing in stroke patients with oropharyngeal dysphagia: A randomized trial. *Journal of Back and Musculoskeletal Rehabilitation*, 33(4), 637–644. <https://doi.org/10.3233/BMR-181133>
- Oh, D.-H., Park, J.-S., & Kim, W.-J. (2017). Effect of neuromuscular electrical stimulation on lip strength and closure function in patients with dysphagia after stroke. *Journal of Physical Therapy Science*, 29(11), 1974–1975. <https://doi.org/10.1589/jpts.29.1974>
- Page, M. J., McKenzie, J. E., Bossuyt, P. M., Boutron, I., Hoffmann, T. C., Mulrow, C. D., Shamseer, L., Tetzlaff, J. M., Akl, E. A., Brennan, S. E., Chou, R., Ghanville, J., Grimshaw, J. M., Hróbjartsson, A., Lalu, M. M., Li, T., Loder, E. W., Mayo-Wilson, E., McDonald, S., McGuinness, L. A., ... Moher, D. (2021). The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ (Clinical research ed.)*, 372, n71. <https://doi.org/10.1136/bmj.n71>
- Park, J.-S., Oh, D.-H., Hwang, N.-K., & Lee, J.-H. (2016). Effects of neuromuscular electrical stimulation combined with effortful swallowing on post-stroke oropharyngeal dysphagia: A randomised controlled trial. *Journal of Oral Rehabilitation*, 43(6), 426–434. <https://doi.org/10.1111/joor.12390>
- Safi, M. F., Wright-Harp, W., Lucker, J. R., Payne, J. C., & Harris, O. (2018). Effect of Neuromuscular Electrical Stimulation on Labial and Lingual Weakness: A Single-Case Preexperimental Study. *Topics in Geriatric Rehabilitation*, 34(2), 145–154. <https://doi.org/10.1097/TGR.000000000000185>
- Sellars, C., Hughes, T., & Langhorne, P. (2005). Speech and language therapy for dysarthria due to non-progressive brain damage. *Em The Cochrane Collaboration (Org.)*, *Cochrane Database of Systematic Reviews* (p. CD002088.pub2). John Wiley & Sons, Ltd. <https://doi.org/10.1002/14651858.CD002088.pub2>
- Sterne, J. A., Hernán, M. A., Reeves, B. C., Savović, J., Berkman, N. D., Viswanathan, M., Henry, D., Altman, D. G., Ansari, M. T., Boutron, I., Carpenter, J. R., Chan,

- A.-W., Churchill, R., Deeks, J. J., Hróbjartsson, A., Kirkham, J., Jüni, P., Loke, Y. K., Pigott, T. D., ... Higgins, J. P. (2016). ROBINS-I: A tool for assessing risk of bias in non-randomised studies of interventions. *BMJ*, i4919. <https://doi.org/10.1136/bmj.i4919>
- Wang, Y., Chung, Y., Chen, H., Chi, L., & Lu, H. (2010). Voice Acoustic Analysis of Taiwanese Adults with Dysarthria Following Stroke. *Canadian Acoustics*, 38, 142-143.
- Wray, F., Clarke, D., & Forster, A. (2019). How do stroke survivors with communication difficulties manage life after stroke in the first year? A qualitative study. *International Journal of Language & Communication Disorders*, 54(5), 814–827. <https://doi.org/10.1111/1460-6984.12487>